

ON THE RIEMANN EXTENSION OF THE GÖDEL SPACE-TIME METRIC

Valery Dryuma

Institute of Mathematics and Informatics AS Moldova, Kishinev

E-mail: valery@dryuma.com; cainar@mail.md dryuma@math.md

Abstract

Some properties of the Gödel space-metric and its Riemann extension are studied.

SATOR AREPO TENET OPERA ROTAS

1 Introduction

The notion of the Riemann extension of nonriemannian spaces was introduced first in ([1]). Main idea of this theory is application of the methods of Riemann geometry for studying of the properties of nonriemanniann spaces.

For example the system differential equations in form

$$\frac{d^2 x^k}{ds^2} + \Pi_{ij}^k \frac{dx^i}{ds} \frac{dx^j}{ds} = 0 \quad (1)$$

with arbitrary coefficients $\Pi_{ij}^k(x^l)$ can be considered as the system of geodesic equations of affinely connected space with local coordinates x^k .

For the n-dimensional Riemannian spaces with the metrics

$${}^n ds^2 = g_{ij} dx^i dx^j$$

the system of geodesic equations looks similar but the coefficients $\Pi_{ij}^k(x^l)$ now have very special form and depends from the choice of the metric g_{ij} .

$$\Pi_{kl}^i = \Gamma_{kl}^i = \frac{1}{2} g^{im} (g_{mk,l} + g_{ml,k} - g_{kl,m})$$

In order that methods of Riemann geometry can be applied for studying of the properties of the spaces with equations (1) the construction of 2n-dimensional extension of the space with local coordinates x^i was introduced .

The metric of extended space constructs with help of coefficients of equation (1) and looks as follows

$${}^{2n} ds^2 = -2\Pi_{ij}^k(x^l) \Psi_k dx^i dx^j + 2d\Psi_k dx^k \quad (2)$$

where Ψ_k are the coordinates of additional space.

The important property of such type metric is that the geodesic equations of metric (2) decomposes into two parts

$$\ddot{x}^k + \Gamma_{ij}^k \dot{x}^i \dot{x}^j = 0, \quad (3)$$

and

$$\frac{\delta^2 \Psi_k}{ds^2} + R_{kji}^l \dot{x}^j \dot{x}^i \Psi_l = 0, \quad (4)$$

where

$$\frac{\delta \Psi_k}{ds} = \frac{d\Psi_k}{ds} - \Gamma_{jk}^l \Psi_l \frac{dx^j}{ds}.$$

The first part (3) of the full system is the system of equations for geodesics of basic space with local coordinates x^i and it does not contain the coordinates Ψ_k .

The second part (4) of system of geodesic equations has the form of linear 4×4 matrix system of second order ODE's for coordinates Ψ_k

$$\frac{d^2 \vec{\Psi}}{ds^2} + A(s) \frac{d\vec{\Psi}}{ds} + B(s) \vec{\Psi} = 0 \quad (5)$$

with the matrix

$$A = A(x^i(s), \dot{x}^i(s)), \quad B = B(x^i(s), \dot{x}^i(s)).$$

From this point of view we have the case of geodesic extension of the basic space (x^i) .

It is important to note that the geometry of extended space is connected with geometry of basic space.

For example the property of the space to be a Ricci-flat

$$R_{ij} = 0,$$

or

$$R_{ij;k} + R_{ki;j} + R_{jk;i} = 0,$$

or symmetrical

$$R_{ijkl;m} = 0$$

keeps also for the extended space.

This fact gives us the possibility to use the linear system of equation (5) for studying of the properties of basic space.

In particular the invariants of 4×4 matrix-function

$$E = B - \frac{1}{2} \frac{dA}{ds} - \frac{1}{4} A^2$$

under change of the coordinates Ψ_k can be of used for that.

For example the condition

$$E = B - \frac{1}{2} \frac{dA}{ds} - \frac{1}{4} A^2 = 0$$

for a given system means that it is equivalent to the simplest system

$$\frac{d^2 \vec{\Phi}}{ds^2} = 0$$

and corresponding extended space is a flat space.

Other cases of integrability of the system (5) are connected with a not flat spaces having special form of the curvature tensor.

Remark that for extended spaces all scalar invariants constructed with the help of curvature tensor and its covariant derivatives are vanishing.

The first applications of the notion of extended spaces to the studying of nonlinear second order differential equations and the Einstein spaces were done in the works of author ([2])-([11]).

Here we consider the properties of the Gödel space-time and its Riemann extension.

2 The Gödel space-time metric

The line element of the metric of the Gödel space-time in coordinates x, y, z, t has the form

$$ds^2 = -dt^2 + dx^2 - 2e^{\frac{x}{a}} dt dy - 1/2 e^{2\frac{x}{a}} dy^2 + dz^2. \quad (6)$$

Here the parameter a is the velocity of rotation ([12]).

The geodesic equations of the metric (6) are given by

$$2 \left(\frac{d^2}{ds^2} x(s) \right) a + \left(e^{\frac{x(s)}{a}} \right)^2 \left(\frac{d}{ds} y(s) \right)^2 + 2 e^{\frac{x(s)}{a}} \left(\frac{d}{ds} t(s) \right) \frac{d}{ds} y(s) = 0, \quad (7)$$

$$\left(\frac{d^2}{ds^2} y(s) \right) e^{\frac{x(s)}{a}} a - 2 \left(\frac{d}{ds} t(s) \right) \frac{d}{ds} x(s) = 0, \quad (8)$$

$$\frac{d^2}{ds^2} z(s) = 0, \quad (9)$$

$$\left(\frac{d^2}{ds^2} t(s) \right) a + e^{\frac{x(s)}{a}} \left(\frac{d}{ds} y(s) \right) \frac{d}{ds} x(s) + 2 \left(\frac{d}{ds} t(s) \right) \frac{d}{ds} x(s) = 0. \quad (10)$$

The first integral of geodesics is satisfied the condition

$$\begin{aligned} & - \left(\frac{d}{ds} t(s) \right)^2 + \left(\frac{d}{ds} x(s) \right)^2 - 2 e^{\frac{x(s)}{a}} \left(\frac{d}{ds} t(s) \right) \frac{d}{ds} y(s) - 1/2 e^{2\frac{x(s)}{a}} \left(\frac{d}{ds} y(s) \right)^2 + \\ & + \left(\frac{d}{ds} z(s) \right)^2 - \mu = 0. \end{aligned}$$

The symbols of Christoffel of the metric (6) are

$$\Gamma_{12}^4 = \frac{\exp(x/a)}{2a}, \quad \Gamma_{14}^2 = -\frac{\exp(-x/a)}{a}, \quad \Gamma_{14}^4 = \frac{1}{a}, \quad \Gamma_{22}^1 = \frac{\exp(2x/a)}{2a}, \quad \Gamma_{24}^1 = \frac{\exp(x/a)}{2a}.$$

To find the solutions of the equations of geodesics (7–10) we present the metric (6) in equivalent form ([13])

$$ds^2 = -(dt + \frac{a\sqrt{2}}{y} dx)^2 + \frac{a^2}{y^2} (dx^2 + dy^2) + dz^2. \quad (11)$$

The correspondence between the both forms of the metrics is given by the relations

$$y = a\sqrt{2} \exp(-x/a), \quad x = y.$$

The equations of geodesics of the metric (11) are defined by

$$\left(\frac{d^2}{ds^2} x(s) \right) a + \sqrt{2} \left(\frac{d}{ds} t(s) \right) \frac{d}{ds} y(s) = 0, \quad (12)$$

$$\left(\frac{d^2}{ds^2} y(s) \right) y(s) a - \left(\frac{d}{ds} x(s) \right)^2 a - \sqrt{2} \left(\frac{d}{ds} t(s) \right) \left(\frac{d}{ds} x(s) \right) y(s) - \left(\frac{d}{ds} y(s) \right)^2 a = 0, \quad (13)$$

$$\left(\frac{d^2}{ds^2} t(s) \right) (y(s))^2 - \sqrt{2} a \left(\frac{d}{ds} y(s) \right) \frac{d}{ds} x(s) - 2 \left(\frac{d}{ds} t(s) \right) \left(\frac{d}{ds} y(s) \right) y(s) = 0, \quad (14)$$

$$\frac{d^2}{ds^2} z(s) = 0. \quad (15)$$

The geodesic equations admit the first integral

$$\begin{aligned}\frac{dt}{ds} &= \frac{(-c_2/\sqrt{2} + \sqrt{2}y)}{c_0}, \\ \frac{dx}{ds} &= \frac{y(c_2 - y)}{ac_0}, \\ \frac{dy}{ds} &= \frac{y(x - c_1)}{ac_0}, \\ \frac{dz}{ds} &= \frac{c_3}{c_0},\end{aligned}\tag{16}$$

where c_i, a_0 are the parameters.

Remark 1 In theory of varieties the Chern-Simons characteristic class is constructed from a matrix gauge connection A_{jk}^i as

$$W(A) = \frac{1}{4\pi^2} \int d^3x \epsilon^{ijk} \text{tr} \left(\frac{1}{2} A_i \partial_j A_k + \frac{1}{3} A_i A_j A_k \right).$$

This term can be translated into a three-dimensional geometric quantity by replacing the matrix connection A_{jk}^i with the Christoffel connection Γ_{jk}^i .

For the density of Chern-Simons invariant can be obtained the expression ([14])

$$CS(\Gamma) = \epsilon^{ijk} (\Gamma_{iq}^p \Gamma_{kp;j}^q + \frac{2}{3} \Gamma_{iq}^p \Gamma_{jr}^q \Gamma_{kp}^r).$$

For the metric (11) at the condition $z = \text{const}$

$$ds^2 = -a^2/y^2 dx^2 - 2\sqrt{2}a/y dx dt + a^2/y^2 dy^2 - dt^2$$

we find the quantity

$$CS(\Gamma) = -\frac{\sqrt{2}}{ay^2}.$$

For the spatial metric

$$^3ds^2 = -g_{\alpha\beta} + \frac{g_{0\alpha}g_{0\beta}}{g_{00}}$$

of the metric (11)

$$-ds^2 = \frac{a^2}{y^2} (dx^2 + dy^2) + dz^2$$

the quantity

$$CS(\Gamma) = 0.$$

3 The Riemann extension of the Gödel metric

The Christoffel symbols of the metric (11) are

$$\Gamma_{11}^2 = -\frac{1}{y}, \quad \Gamma_{22}^2 = -\frac{1}{y}, \quad \Gamma_{14}^2 = -\frac{\sqrt{2}}{2a}, \quad \Gamma_{24}^1 = \frac{\sqrt{2}}{2a}, \quad \Gamma_{12}^4 = -\frac{a\sqrt{2}}{2y^2}, \quad \Gamma_{14}^4 = -\frac{1}{y}.$$

Now with help of the formulae (2) we construct the eight-dimensional extension of the metric (11). It has the form

$$^8ds^2 = \frac{2}{y} Q dx^2 + \frac{2a\sqrt{2}}{y^2} V dx dy + \frac{2\sqrt{2}}{a} Q dx dt + \frac{2}{y} Q dy^2 + \left(\frac{4}{y} V - 2\frac{\sqrt{2}}{a} P \right) dy dt +$$

$$+2dx dP + 2dy dQ + 2dz dU + 2dt dV. \quad (17)$$

where (P, Q, U, V) are an additional coordinates.

The Ricci tensor of the four-dimensional Gödel space with the metric (11) or (6) satisfied the condition

$${}^4R_{ik;l} + {}^4R_{li;k} + {}^4R_{kl;i} = 0.$$

This property is valid for the eight-dimensional space in local coordinates (x, y, z, t, P, Q, U, V) with the metric (11)

$${}^8R_{ik;l} + {}^8R_{li;k} + {}^8R_{kl;i} = 0.$$

The full system of geodesic equations for the metric (7) decomposes into two parts.

The first part coincides with the equations (12-15) on the coordinates (x, y, z, t) and second part forms the linear system of equations for coordinates P, Q, U, V .

They are defined as

$$\begin{aligned} \frac{d^2}{ds^2}P(s) = & -\frac{\left(\sqrt{2}a^2\left(\frac{d}{ds}x(s)\right)^2 + 2\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}x(s)\right)y(s)a - \sqrt{2}a^2\left(\frac{d}{ds}y(s)\right)^2\right)V(s)}{(y(s))^3a} \\ & -\frac{\left(2\left(\frac{d}{ds}x(s)\right)(y(s))^2a + \sqrt{2}\left(\frac{d}{ds}t(s)\right)(y(s))^3\right)\frac{d}{ds}Q(s)}{(y(s))^3a} - \frac{\sqrt{2}a\left(\frac{d}{ds}V(s)\right)\frac{d}{ds}y(s)}{(y(s))^2}, \end{aligned} \quad (18)$$

$$\begin{aligned} \frac{d^2}{ds^2}Q(s) = & \frac{\left(-3\left(\frac{d}{ds}x(s)\right)^2y(s)a - 2\sqrt{2}\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}x(s)\right)(y(s))^2 - \left(\frac{d}{ds}y(s)\right)^2y(s)a\right)Q(s)}{(y(s))^3a} + \\ & + \frac{\left(2a\left(\frac{d}{ds}y(s)\right)\left(\frac{d}{ds}x(s)\right)y(s) + 2\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}y(s)\right)(y(s))^2\sqrt{2}\right)P(s)}{(y(s))^3a} + \\ & + \frac{\left(-2\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}y(s)\right)y(s)a - 2\sqrt{2}a^2\left(\frac{d}{ds}y(s)\right)\frac{d}{ds}x(s)V(s) + \sqrt{2}\left(\frac{d}{ds}t(s)\right)\frac{d}{ds}P(s)\right)}{(y(s))^3a} - \\ & - 2\frac{\left(\frac{d}{ds}y(s)\right)\frac{d}{ds}Q(s)}{y(s)} + \frac{\left(-\sqrt{2}a^2\left(\frac{d}{ds}x(s)\right)y(s) - 2\left(\frac{d}{ds}t(s)\right)(y(s))^2a\right)\frac{d}{ds}V(s)}{(y(s))^3a}, \end{aligned} \quad (19)$$

$$\frac{d^2}{ds^2}U(s) = 0, \quad (20)$$

$$\begin{aligned} \frac{d^2}{ds^2}V(s) = & \frac{\left(\left(\frac{d}{ds}x(s)\right)^2a\sqrt{2}y(s) + 2\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}x(s)\right)(y(s))^2 + \sqrt{2}\left(\frac{d}{ds}y(s)\right)^2ay(s)\right)P(s)}{a^2(y(s))^2} + \\ & + \frac{\left(-2\sqrt{2}\left(\frac{d}{ds}t(s)\right)\left(\frac{d}{ds}x(s)\right)y(s)a - 2a^2\left(\frac{d}{ds}x(s)\right)^2\right)V(s)}{a^2(y(s))^2} + \frac{\sqrt{2}\left(\frac{d}{ds}y(s)\right)\frac{d}{ds}P(s)}{a} \\ & - \frac{\sqrt{2}\left(\frac{d}{ds}x(s)\right)\frac{d}{ds}Q(s)}{a} - 2\frac{\left(\frac{d}{ds}y(s)\right)\frac{d}{ds}V(s)}{y(s)} + 2\frac{Q(s)\left(\frac{d}{ds}t(s)\right)\frac{d}{ds}y(s)}{a^2}. \end{aligned} \quad (21)$$

In result we have got a linear matrix-second order ODE for the coordinates U, V, P, Q

$$\frac{d^2\Psi}{ds^2} = A(x, \phi, z, t)\frac{d\Psi}{ds} + B(x, \phi, z, t)\Psi, \quad (22)$$

where

$$\Psi(s) = \begin{pmatrix} P(s) \\ Q(s) \\ U(s) \\ V(s) \end{pmatrix}$$

and A, B are some 4×4 matrix-functions depending from the coordinates $x(s), y(s), z(s), t(s)$ and their derivatives.

Now we shall investigate the properties of the matrix system of equations (18-21).

To integrate this system we use the relation

$$\dot{x}(s)P(s) + \dot{y}(s)Q(s) + \dot{z}(s)U(s) + \dot{t}(s)V(s) - \frac{s}{2} - \mu = 0, \quad (23)$$

which is valid for the every Riemann extensions of affinely connected space and which is consequence of the well known first integral of geodesic equations

$$g_{ik}\dot{x}^i\dot{x}^k = \nu$$

of arbitrary Riemann space.

Using the expressions for the first integrals of geodesic (16) and $U(s) = \alpha s + \beta$ from the equation (20) the system of equations (18-21) may be simplified.

In result we get the system of equations for additional coordinates

$$\begin{aligned} \frac{d^2}{ds^2}P(s) &= \frac{(\sqrt{2}c\theta ac1 - \sqrt{2}c\theta ax(s)) \frac{d}{ds}V(s)}{y(s)c\theta^2a} + \\ &+ \frac{(-\sqrt{2}c2y(s) - 2\sqrt{2}x(s)c1 + \sqrt{2}c1^2 + \sqrt{2}(y(s))^2 + \sqrt{2}(x(s))^2)V(s)}{y(s)c\theta^2a} - \frac{(\frac{d}{ds}Q(s))c2}{c\theta a}, \end{aligned} \quad (24)$$

$$\begin{aligned} \frac{d^2}{ds^2}Q(s) &= -\frac{(2(y(s))^2c1 - 2(y(s))^2x(s))P(s)}{y(s)a^2c\theta^2} - \\ &- \frac{(y(s)(x(s))^2 - (y(s))^3 + y(s)c1^2 + y(s)c2^2 - 2y(s)x(s)c1)Q(s)}{y(s)a^2c\theta^2} - \\ &- \frac{(2ac\theta y(s)x(s) - 2ac\theta y(s)c1)\frac{d}{ds}Q(s)}{y(s)a^2c\theta^2} - \frac{(ac\theta y(s)c2 - 2ac\theta(y(s))^2)\frac{d}{ds}P(s)}{y(s)a^2c\theta^2} - \\ &- \frac{\sqrt{2}\frac{d}{ds}V(s)}{c\theta} - \frac{(\sqrt{2}c2ax(s) - \sqrt{2}c2ac1)V(s)}{y(s)a^2c\theta^2}, \end{aligned} \quad (25)$$

$$\begin{aligned} \frac{d^2}{ds^2}V(s) &= \frac{((y(s))^2\sqrt{2}c2 - (y(s))^3\sqrt{2} + y(s)\sqrt{2}(x(s))^2 - 2y(s)\sqrt{2}x(s)c1 + y(s)\sqrt{2}c1^2)P(s)}{c\theta^2a^3} + \\ &+ \frac{(-\sqrt{2}y(s)x(s)c2 + \sqrt{2}y(s)c1c2 + 2\sqrt{2}(y(s))^2x(s) - 2\sqrt{2}(y(s))^2c1)Q(s)}{c\theta^2a^3} + \\ &+ \frac{(-\sqrt{2}y(s)ac\theta c2 + \sqrt{2}(y(s))^2ac\theta)\frac{d}{ds}Q(s)}{c\theta^2a^3} + \frac{(\sqrt{2}y(s)ac\theta x(s) - \sqrt{2}y(s)ac\theta c1)\frac{d}{ds}P(s)}{c\theta^2a^3} + \end{aligned}$$

$$+ \frac{(-2a^2c\theta x(s) + 2a^2c\theta c1) \frac{d}{ds}V(s)}{c\theta^2a^3} + \frac{(-2ac2y(s) + 2a(y(s))^2) V(s)}{c\theta^2a^3}. \quad (26)$$

The relation (23) in this case takes a form

$$\begin{aligned} -1/2 \frac{(-2\alpha c3a + c\theta a)s}{c\theta a} - 1/2 \frac{(c2\sqrt{2}a - 2y(s)\sqrt{2}a) V(s)}{c\theta a} - 1/2 \frac{(2c1y(s) - 2x(s)y(s)) Q(s)}{c\theta a} - \\ - 1/2 \frac{(2(y(s))^2 - 2c2y(s)) P(s)}{c\theta a} - \nu = 0 \end{aligned} \quad (27)$$

and the system from three equations (24-26) can be reduced to the system of two coupled equations.

As example the substitution of the expression

$$\begin{aligned} Q(s) = 1/2 \frac{(-2\alpha c3a + c\theta a)s}{y(s)(x(s) - c1)} + 1/2 \frac{(c2\sqrt{2}a - 2y(s)\sqrt{2}a) V(s)}{y(s)(x(s) - c1)} + 1/2 \frac{(2(y(s))^2 - 2c2y(s)) P(s)}{y(s)(x(s) - c1)} + \\ + \frac{\nu c\theta a}{y(s)(x(s) - c1)} \end{aligned}$$

into the equation for $Q(s)$ give us the identity and in result our system takes the form

$$\begin{aligned} \frac{d^2}{ds^2}P(s) = - \frac{(-c2 + y(s)) c2 \frac{d}{ds}P(s)}{c\theta a(x(s) - c1)} - 1/2 \frac{(2(x(s))^2 - 4x(s)c1 - 2c2y(s) + c2^2 + 2c1^2) \sqrt{2} \frac{d}{ds}V(s)}{c\theta (x(s) - c1)y(s)} - \\ - \frac{c2y(s) ((y(s))^2 + c2^2 - 2c2y(s) + c1^2 - 2x(s)c1 + (x(s))^2) P(s)}{a^2c\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} + L(s)V(s) - \\ - 1/2 \frac{(-2\alpha c3ac2 + c\theta ac2) sy(s)}{a^2c\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} - 1/2 \frac{(-c\theta ac2^2 + 2\alpha c3ac2^2) s}{a^2c\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} - \\ - 1/2 \frac{((2\alpha c3ac2 - c\theta ac2)(x(s))^2 + (2c1c\theta ac2 - 4\alpha c3c1ac2)x(s) + 2\alpha c3c1^2ac2 - c1^2c\theta ac2) s}{a^2c\theta^2 y(s) ((x(s))^2 - 2x(s)c1 + c1^2)}, \end{aligned} \quad (28)$$

where

$$\begin{aligned} L(s) = \frac{(y(s))^2 \sqrt{2}c2}{ac\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} + 1/2 \frac{\sqrt{2} (2(x(s))^2 - 3c2^2 - 4x(s)c1 + 2c1^2) y(s)}{ac\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} + \\ + 1/2 \frac{\sqrt{2} (-2c2(x(s))^2 - 2c1^2c2 + c2^3 + 4c2c1x(s))}{ac\theta^2 ((x(s))^2 - 2x(s)c1 + c1^2)} + \\ + 1/2 \frac{\sqrt{2} (2(x(s))^4 - 8(x(s))^3c1 + (c2^2 + 12c1^2)(x(s))^2 + (-2c2^2c1 - 8c1^3)x(s) + c1^2c2^2 + 2c1^4)}{ac\theta^2 y(s) ((x(s))^2 - 2x(s)c1 + c1^2)}, \end{aligned}$$

and

$$\frac{d^2}{ds^2}V(s) = - \frac{(2(x(s))^2 - 4x(s)c1 - 3c2y(s) + 2(y(s))^2 + 2c1^2 + c2^2) \frac{d}{ds}V(s)}{ac\theta (x(s) - c1)} +$$

$$\begin{aligned}
& + \frac{\left((y(s))^2 + c^2 - 2c^2 y(s) + c^2 - 2x(s)c + (x(s))^2 \right) \sqrt{2} y(s) \frac{d}{ds} P(s)}{c \theta a^2 (x(s) - c)} + M(s)V(s) + N(s)P(s) + \\
& + 1/2 \frac{\sqrt{2} (c\theta - 2\alpha c^3) s (y(s))^3}{a^2 c \theta^2 (x(s) - c^2)} + 1/2 \frac{\sqrt{2} (-2c\theta c^2 + 4\alpha c^3 c^2) s (y(s))^2}{a^2 c \theta^2 (x(s) - c^2)} + \\
& + 1/2 \frac{\sqrt{2} \left(-2\alpha c^3 c^2 + c^2 c\theta + c^2 c\theta + (x(s))^2 c\theta - 2x(s)c\theta c - 2\alpha c^3 c^2 + 4\alpha c^3 x(s)c - 2\alpha c^3 (x(s))^2 \right) s y(s)}{a^2 c \theta^2 (x(s) - c^2)},
\end{aligned} \tag{29}$$

where

$$\begin{aligned}
M(s) &= -2 \frac{(y(s))^4}{a^2 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} + 5 \frac{(y(s))^3 c^2}{a^2 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} - \\
& - \frac{\left(-4x(s)c + 2(x(s))^2 + 4c^2 + 2c^2 \right) (y(s))^2}{a^2 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} - \frac{\left(-c^2 (x(s))^2 - c^2 c^2 + 2c^2 c x(s) - c^2 c^3 \right) y(s)}{a^2 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)}, \\
N(s) &= \frac{\sqrt{2} (y(s))^5}{a^3 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} - 3 \frac{\sqrt{2} (y(s))^4 c^2}{a^3 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} + \\
& + \frac{\sqrt{2} \left(2(x(s))^2 + 3c^2 + 2c^2 - 4x(s)c \right) (y(s))^3}{a^3 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} + \frac{\sqrt{2} \left(-3c^2 (x(s))^2 - c^2 c^3 - 3c^2 c^2 + 6c^2 c x(s) \right) (y(s))^2}{a^3 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)} + \\
& + \frac{\sqrt{2} \left((x(s))^4 - 4(x(s))^3 c + (6c^2 + c^2) (x(s))^2 + (-2c^2 c - 4c^3) x(s) + c^4 + c^2 c^2 \right) y(s)}{a^3 c \theta^2 \left((x(s))^2 - 2x(s)c + c^2 \right)}.
\end{aligned}$$

The expressions for functions $x(s)$ and $y(s)$ are dependent from the choice of parameters and can be defined from the equations (16).

The integration of the equations (28-29) for the additional coordinates $P(s)$, $Q(s)$ is reduced to investigation of the 2×2 system of second order ODE's with variable coefficients.

Remark that the matrix E and its properties play important role in analysis of such type of the system of equations.

In result we get the correspondence between the geodesic in the x, y, x, t -space and the geodesic in the space with local coordinates P, Q, U, V (partner space).

The studying of such type of correspondence may be useful from various point of view.

4 Translation surfaces of the Gödel spaces

Now we discuss the properties of translation surfaces of the Gödel spaces.

According with definition ([15]) translation surfaces in arbitrary Riemannian space are defined by the systems of equations for local coordinates $x^i(u, v)$ of the space

$$\frac{\partial x^i(u, v)}{\partial u \partial v} + \Gamma_{jk}^i(x^m) \frac{\partial x^j(u, v)}{\partial u} \frac{\partial x^k(u, v)}{\partial v} = 0, \tag{30}$$

where Γ_{jk}^i are the Christoffel coefficients.

In the case of the Gödel metric (6) we get the equations

$$\frac{\partial^2}{\partial u \partial v} x(u, v) + 1/2 \frac{\sqrt{2} \left(\left(\frac{\partial}{\partial u} y(u, v) \right) \frac{\partial}{\partial v} t(u, v) + \left(\frac{\partial}{\partial u} t(u, v) \right) \frac{\partial}{\partial v} y(u, v) \right)}{a} = 0, \quad (31)$$

$$\begin{aligned} & \frac{\partial^2}{\partial u \partial v} y(u, v) - 1/2 \frac{\sqrt{2} \left(\left(\frac{\partial}{\partial v} x(u, v) \right) \frac{\partial}{\partial u} t(u, v) - \left(\frac{\partial}{\partial v} t(u, v) \right) \frac{\partial}{\partial u} x(u, v) \right)}{a} \\ & - \frac{\left(\frac{\partial}{\partial u} x(u, v) \right) \frac{\partial}{\partial v} x(u, v) + \left(\frac{\partial}{\partial u} y(u, v) \right) \frac{\partial}{\partial v} y(u, v)}{y(u, v)} = 0, \end{aligned} \quad (32)$$

$$\begin{aligned} & \frac{\partial^2}{\partial u \partial v} t(u, v) - \frac{\left(\frac{\partial}{\partial u} y(u, v) \right) \frac{\partial}{\partial v} t(u, v) + \left(\frac{\partial}{\partial u} t(u, v) \right) \frac{\partial}{\partial v} y(u, v)}{y(u, v)} - \\ & - \frac{1/2 \sqrt{2} a \left(\frac{\partial}{\partial u} x(u, v) \right) \frac{\partial}{\partial v} y(u, v) + 1/2 \sqrt{2} a \left(\frac{\partial}{\partial u} y(u, v) \right) \frac{\partial}{\partial v} x(u, v)}{(y(u, v))^2} = 0 \end{aligned} \quad (33)$$

$$\frac{\partial^2}{\partial u \partial v} z(u, v) = 0 \quad (34)$$

Full integration of this nonlinear system of equations is a difficult problem.

To cite one example.

With this aim we present our system of equations in a new coordinates $u = r + s, v = r - s$.

It takes the form

$$\begin{aligned} & 2 \left(1/4 \frac{\partial^2}{\partial r^2} x(r, s) - 1/4 \frac{\partial^2}{\partial s^2} x(r, s) \right) a + \sqrt{2} \left(1/2 \frac{\partial}{\partial r} y(r, s) + 1/2 \frac{\partial}{\partial s} y(r, s) \right) \left(1/2 \frac{\partial}{\partial r} t(r, s) - 1/2 \frac{\partial}{\partial s} t(r, s) \right) + \\ & + \sqrt{2} \left(1/2 \frac{\partial}{\partial r} t(r, s) + 1/2 \frac{\partial}{\partial s} t(r, s) \right) \left(1/2 \frac{\partial}{\partial r} y(r, s) - 1/2 \frac{\partial}{\partial s} y(r, s) \right) = 0, \\ & -2 \left(1/4 \frac{\partial^2}{\partial r^2} y(r, s) - 1/4 \frac{\partial^2}{\partial s^2} y(r, s) \right) y(r, s) a + \\ & + 2 \left(1/2 \frac{\partial}{\partial r} x(r, s) + 1/2 \frac{\partial}{\partial s} x(r, s) \right) \left(1/2 \frac{\partial}{\partial r} x(r, s) - 1/2 \frac{\partial}{\partial s} x(r, s) \right) a + \\ & + \sqrt{2} \left(1/2 \frac{\partial}{\partial r} x(r, s) + 1/2 \frac{\partial}{\partial s} x(r, s) \right) \left(1/2 \frac{\partial}{\partial r} t(r, s) - 1/2 \frac{\partial}{\partial s} t(r, s) \right) y(r, s) + \\ & + 2 \left(1/2 \frac{\partial}{\partial r} y(r, s) + 1/2 \frac{\partial}{\partial s} y(r, s) \right) \left(1/2 \frac{\partial}{\partial r} y(r, s) - 1/2 \frac{\partial}{\partial s} y(r, s) \right) a + \\ & + \sqrt{2} \left(1/2 \frac{\partial}{\partial r} t(r, s) + 1/2 \frac{\partial}{\partial s} t(r, s) \right) \left(1/2 \frac{\partial}{\partial r} x(r, s) - 1/2 \frac{\partial}{\partial s} x(r, s) \right) y(r, s) = 0, \\ & -2 \left(1/4 \frac{\partial^2}{\partial r^2} t(r, s) - 1/4 \frac{\partial^2}{\partial s^2} t(r, s) \right) (y(r, s))^2 + \\ & + \sqrt{2} a \left(1/2 \frac{\partial}{\partial r} x(r, s) + 1/2 \frac{\partial}{\partial s} x(r, s) \right) \left(1/2 \frac{\partial}{\partial r} y(r, s) - 1/2 \frac{\partial}{\partial s} y(r, s) \right) + \end{aligned}$$

$$\begin{aligned}
& +\sqrt{2}a \left(1/2 \frac{\partial}{\partial r} y(r, s) + 1/2 \frac{\partial}{\partial s} y(r, s) \right) \left(1/2 \frac{\partial}{\partial r} x(r, s) - 1/2 \frac{\partial}{\partial s} x(r, s) \right) + \\
& +2 \left(1/2 \frac{\partial}{\partial r} y(r, s) + 1/2 \frac{\partial}{\partial s} y(r, s) \right) \left(1/2 \frac{\partial}{\partial r} t(r, s) - 1/2 \frac{\partial}{\partial s} t(r, s) \right) y(r, s) + \\
& +2 \left(1/2 \frac{\partial}{\partial r} t(r, s) + 1/2 \frac{\partial}{\partial s} t(r, s) \right) \left(1/2 \frac{\partial}{\partial r} y(r, s) - 1/2 \frac{\partial}{\partial s} y(r, s) \right) y(r, s) = 0.
\end{aligned}$$

The solution of this system of equations we shall seek in form

$$y(r, s) = B(r), \quad t(r, s) = C(r) - s, \quad x(r, s) = s + A(r),$$

where $B(r), C(r), A(r)$ are some unknown functions.

In result our system takes the form

$$\begin{aligned}
& \left(\frac{d^2}{dr^2} C(r) \right) (B(r))^2 - \sqrt{2}a \left(\frac{d}{dr} B(r) \right) \frac{d}{dr} A(r) - 2 \left(\frac{d}{dr} B(r) \right) B(r) \frac{d}{dr} C(r) = 0, \\
& \left(\frac{d^2}{dr^2} B(r) \right) B(r)a - a \left(\frac{d}{dr} A(r) \right)^2 + a - \sqrt{2}B(r) \left(\frac{d}{dr} A(r) \right) \frac{d}{dr} C(r) - \sqrt{2}B(r) - \left(\frac{d}{dr} B(r) \right)^2 a = 0, \\
& \left(\frac{d^2}{dr^2} A(r) \right) a + \sqrt{2} \left(\frac{d}{dr} B(r) \right) \frac{d}{dr} C(r) = 0.
\end{aligned}$$

Using the first integral

$$\frac{d}{dr} C(r) = -\frac{\sqrt{2}a \frac{d}{dr} A(r)}{B(r)} + \alpha,$$

the system can be written in form

$$\begin{aligned}
& \left(\frac{d^2}{dr^2} B(r) \right) B(r)a + a \left(\frac{d}{dr} A(r) \right)^2 + a - \sqrt{2} \left(\frac{d}{dr} A(r) \right) \alpha B(r) - \sqrt{2}B(r) - \left(\frac{d}{dr} B(r) \right)^2 a = 0, \\
& \left(\frac{d^2}{dr^2} A(r) \right) aB(r) - 2 \left(\frac{d}{dr} B(r) \right) a \frac{d}{dr} A(r) + \sqrt{2} \left(\frac{d}{dr} B(r) \right) \alpha B(r) = 0.
\end{aligned}$$

Integration of the last equation give us the expression for the function $A(r)$

$$A(r) = \int \frac{B(r) (\sqrt{2}\alpha + B(r) C_1 a)}{a} dr + C_2$$

with parameters C_1, C_2 and α .

After substitution the expression for $A(r)$ to the first equation we get the equation

$$\left(\frac{d^2}{dr^2} B(r) \right) B(r)a + (B(r))^3 \sqrt{2}\alpha C_1 + (B(r))^4 C_1^2 a + a - \sqrt{2}B(r) - \left(\frac{d}{dr} B(r) \right)^2 a = 0$$

for the function $B(r)$.

Remark that this equation can be written in form

$$\frac{d^2}{dr^2} E(r) = \frac{\sqrt{2}e^{-E(r)}}{a} - C_1^2 e^{2E(r)} - \frac{\sqrt{2}\alpha C_1 e^{E(r)}}{a} - e^{-2E(r)}$$

after the change of variable

$$B(r) = \exp(E(r)).$$

With the help of its solutions the examples of the translation surfaces of the Gödel space (6) can be constructed.

They can be presented in form

$$t + x = A(r) + C(r), \quad y = B(r),$$

or

$$t(x, y) = x + \phi(y)$$

with some function $\phi(y)$.

Detail consideration of the properties of this type of translation surfaces, their intrinsic geometry and characteristic lines will be done in a following publications of author.

Remark that with the help of the translation surfaces can be investigated the properties of closed trajectories of the Gödel space.

Let us consider the eight-dimensional extension of the Gödel space with the metric (17).

Translation surfaces in this case are determined by the equations (28)-(31) for coordinates x, y, z, t and by the linear system of equations on coordinates P, Q, U, V

$$\begin{aligned} & \frac{\partial^2}{\partial u \partial v} P(u, v) - \\ & -1/2 \frac{(-2\sqrt{2}a^2 (\frac{\partial}{\partial u} x) \frac{\partial}{\partial v} x - 2 (\frac{\partial}{\partial u} x) (\frac{\partial}{\partial v} t) ya + 2\sqrt{2}a^2 (\frac{\partial}{\partial u} y) \frac{\partial}{\partial v} y - 2 (\frac{\partial}{\partial u} t) (\frac{\partial}{\partial v} x) ya) V(u, v)}{y^3 a} \\ & -1/2 \frac{(-2 (\frac{\partial}{\partial u} x(u, v)) y^2 a - \sqrt{2} (\frac{\partial}{\partial u} t(u, v)) y^3) \frac{\partial}{\partial v} Q(u, v)}{y^3 a} - \\ & -1/2 \frac{(-\sqrt{2} (\frac{\partial}{\partial v} t(u, v)) y^3 - 2 (\frac{\partial}{\partial v} x(u, v)) y^2 a) \frac{\partial}{\partial u} Q(u, v)}{y^3 a} + \\ & +1/2 \frac{\sqrt{2} a (\frac{\partial}{\partial u} y(u, v)) \frac{\partial}{\partial v} V(u, v)}{y^2} + 1/2 \frac{\sqrt{2} a (\frac{\partial}{\partial v} y(u, v)) \frac{\partial}{\partial u} V(u, v)}{y^2} = 0, \end{aligned} \quad (35)$$

$$\begin{aligned} & \frac{\partial^2}{\partial u \partial v} Q(u, v) + \\ & +1/2 \frac{(6 (\frac{\partial}{\partial u} x) (\frac{\partial}{\partial v} x) ya + 2\sqrt{2} (\frac{\partial}{\partial u} x) (\frac{\partial}{\partial v} t) y^2 + 2\sqrt{2} (\frac{\partial}{\partial u} t) (\frac{\partial}{\partial v} x) y^2 + 2 (\frac{\partial}{\partial u} y) (\frac{\partial}{\partial v} y) ya) Q(u, v)}{y^3 a} + \\ & +1/2 \frac{(-2a (\frac{\partial}{\partial u} x) (\frac{\partial}{\partial v} y) y - 2 (\frac{\partial}{\partial u} y) (\frac{\partial}{\partial v} t) y^2 \sqrt{2} - 2 (\frac{\partial}{\partial u} t) (\frac{\partial}{\partial v} y) y^2 \sqrt{2} - 2a (\frac{\partial}{\partial u} y) (\frac{\partial}{\partial v} x) y) P(u, v)}{y^3 a} + \\ & +1/2 \frac{(2\sqrt{2}a^2 (\frac{\partial}{\partial u} x) \frac{\partial}{\partial v} y + 2 (\frac{\partial}{\partial u} y) (\frac{\partial}{\partial v} t) ya + 2\sqrt{2}a^2 (\frac{\partial}{\partial u} y) \frac{\partial}{\partial v} x + 2 (\frac{\partial}{\partial u} t) (\frac{\partial}{\partial v} y) ya) V(u, v)}{y^3 a} + \\ & + \frac{(\frac{\partial}{\partial u} y(u, v)) \frac{\partial}{\partial v} Q(u, v)}{y} + \frac{(\frac{\partial}{\partial v} y(u, v)) \frac{\partial}{\partial u} Q(u, v)}{y} + 1/2 \frac{(2 (\frac{\partial}{\partial u} t(u, v)) y^2 a + \sqrt{2}a^2 (\frac{\partial}{\partial u} x(u, v)) y) \frac{\partial}{\partial v} V(u, v)}{y^3 a} + \\ & +1/2 \frac{(\sqrt{2}a^2 (\frac{\partial}{\partial v} x(u, v)) y + 2 (\frac{\partial}{\partial v} t(u, v)) y^2 a) \frac{\partial}{\partial u} V(u, v)}{y^3 a} + \\ & +1/2 \frac{-\sqrt{2} (\frac{\partial}{\partial u} t(u, v)) (\frac{\partial}{\partial v} P(u, v)) y^3 - \sqrt{2} (\frac{\partial}{\partial u} P(u, v)) (\frac{\partial}{\partial v} t(u, v)) y^3}{y^3 a} = 0, \end{aligned} \quad (36)$$

$$\frac{\partial^2}{\partial u \partial v} V(u, v) +$$

$$\begin{aligned}
& +1/2 \frac{(-2 \left(\frac{\partial}{\partial u} y(u, v)\right) \left(\frac{\partial}{\partial v} t(u, v)\right) y^2 - 2 \left(\frac{\partial}{\partial u} t(u, v)\right) \left(\frac{\partial}{\partial v} y(u, v)\right) y^2) Q(u, v)}{a^2 y^2} + \\
& +1/2 \frac{(-2 \left(\frac{\partial}{\partial u} x\right) \left(\frac{\partial}{\partial v} t\right) y^2 - 2 \left(\frac{\partial}{\partial u} x\right) \left(\frac{\partial}{\partial v} x\right) a \sqrt{2} y - 2 \sqrt{2} \left(\frac{\partial}{\partial u} y\right) \left(\frac{\partial}{\partial v} y\right) a y - 2 \left(\frac{\partial}{\partial u} t\right) \left(\frac{\partial}{\partial v} x\right) y^2) P(u, v)}{a^2 y^2} + \\
& +1/2 \frac{(4 \left(\frac{\partial}{\partial u} x\right) \left(\frac{\partial}{\partial v} x\right) a^2 + 2 \sqrt{2} \left(\frac{\partial}{\partial u} x\right) \left(\frac{\partial}{\partial v} t\right) y a + 2 \sqrt{2} \left(\frac{\partial}{\partial u} t\right) \left(\frac{\partial}{\partial v} x\right) y a) V(u, v)}{a^2 y^2} + \\
& +1/2 \frac{\sqrt{2} \left(\frac{\partial}{\partial u} x(u, v)\right) \frac{\partial}{\partial v} Q(u, v)}{a} + \\
& +1/2 \frac{\sqrt{2} \left(\frac{\partial}{\partial u} Q(u, v)\right) \frac{\partial}{\partial v} x(u, v)}{a} + \frac{\left(\frac{\partial}{\partial u} y(u, v)\right) \frac{\partial}{\partial v} V(u, v)}{y} + \frac{\left(\frac{\partial}{\partial u} V(u, v)\right) \frac{\partial}{\partial v} y(u, v)}{y} + \\
& +1/2 \frac{-\sqrt{2} \left(\frac{\partial}{\partial u} P(u, v)\right) \left(\frac{\partial}{\partial v} y(u, v)\right) a y^2 - \sqrt{2} \left(\frac{\partial}{\partial u} y(u, v)\right) \left(\frac{\partial}{\partial v} P(u, v)\right) a y^2}{a^2 y^2} = 0 \tag{37}
\end{aligned}$$

$$\frac{\partial^2}{\partial u \partial v} U(u, v) = 0. \tag{38}$$

The system of linear equations (32)-(35) is the matrix analog of the Laplace equation

$$\frac{\partial^2 \Psi(u, v)}{\partial u \partial v} + A(u, v) \frac{\partial \Psi(u, v)}{\partial u} + B(u, v) \frac{\partial \Psi(u, v)}{\partial v} + C(u, v) \Psi(u, v) = 0, \tag{39}$$

where

$$\Psi(u, v) = \begin{pmatrix} P(u, v) \\ Q(u, v) \\ U(u, v) \\ V(u, v) \end{pmatrix}$$

is a vector-function, and $A(u, v), B(u, v), C(u, v)$ are the matrices depending from the variables (u, v) .

For integration of such type equation can be used the matrix generalization of the Darboux Invariants ([16]).

Remark 2 We are reminded basic facts on integration of the matrix Laplace-equation.

The system (39) can be presented in forms

$$(\partial_u + B)(\partial_v + A)\Psi - H\Psi = 0,$$

or

$$(\partial_v + A)(\partial_u + B)\Psi - K\Psi = 0,$$

where

$$H = \frac{\partial A}{\partial u} + BA - C, \quad K = \frac{\partial B}{\partial v} + AB - C,$$

are the Darboux invariants of the system.

In the case $K = 0$ or $H = 0$ the system can be integrated.

If the $K \neq 0$ and $H \neq 0$ the system may presented in similar form for the functions

$$\Psi_1 = \frac{\partial \Psi}{\partial y} + A\Psi.$$

or

$$\Psi_{-1} = \frac{\partial \Psi}{\partial x} + B\Psi.$$

In the first case one get

$$\frac{\partial^2 \Psi_1(u, v)}{\partial u \partial v} + A_1(u, v) \frac{\partial \Psi_1(u, v)}{\partial u} + B_1(u, v) \frac{\partial \Psi_1(u, v)}{\partial v} + C_1(u, v) \Psi_1(u, v) = 0,$$

where

$$A_1 = HAH^{-1} - H_vH^{-1}, \quad B_1 = B, \quad C_1 = B_v - H + (HAH^{-1} - H_vH^{-1})B.$$

The invariants H_1 and K_1 for this equation are

$$H_1 = H - B_y + (HAH^{-1} - H_yH^{-1})_u + B(HAH^{-1} - H_vH^{-1}) - (HAH^{-1} - H_yH^{-1})B,$$

$$K_1 = H.$$

In the case $H_1 = 0$ the system can be integrated.

In the second case we get the equation for the function Ψ_{-1}

$$\frac{\partial^2 \Psi_{-1}(u, v)}{\partial u \partial v} + A_{-1}(u, v) \frac{\partial \Psi_{-1}(u, v)}{\partial u} + B_{-1}(u, v) \frac{\partial \Psi_{-1}(u, v)}{\partial v} + C_{-1}(u, v) \Psi_{-1}(u, v) = 0,$$

where

$$B_{-1} = KBK^{-1} - K_uK^{-1}, \quad A_{-1} = A, \quad C_{-1} = A_u - K + (KBK^{-1} - K_uK^{-1})A.$$

The invariants H_{-1} and K_{-1} for this equation are

$$K_{-1} = K - A_u + (KBK^{-1} - K_uK^{-1})_v + A(KBK^{-1} - K_uK^{-1}) - (KBK^{-1} - K_uK^{-1})A,$$

$$H_{-1} = K$$

and at the condition $K_{-1} = 0$ the system is also integrable.

To integrate the system of equations (39) in explicit form it is necessary to use the expressions for coordinates $x(u, v), y(u, v), z(u, v), t(u, v)$ of translation surfaces of the basic space.

The properties of the invariants H and K also may be important for classifications of translation surfaces of the basic and extended Gödel space.

5 On the spectrum of the Gödel space-time metric

In this section the spectrum λ of de Rham operator

$$\Delta = g^{ij} \nabla_i \nabla_j - Ricci,$$

defined on a four-dimensional riemannian manifold and acting on 1-forms

$$\omega = A_i(x, y, z) dx^i = u(x, y, z, t) dx + v(x, y, z, t) dy + p(x, y, z, t) dz + q(x, y, z, t) dt$$

will be calculated.

The problem is reduced to the solution of the system of equations

$$g^{ij} \nabla_i \nabla_j A_k - R_k^l A_l - \mu^2 A_k = 0, \tag{40}$$

where ∇_k is a symbol of covariant derivative and the R_j^i is the Ricci tensor of the metric g^{ij} of the Gödel space-time.

We use the Gödel space-time metric in form (11) and for simplicity sake the components of the 1-form ω will be presented as

$$A_k = [0, v(y, t), 0, q(y, t)].$$

As this takes place the system (40) looks as

$$-\frac{\partial}{\partial t} v(y, t) + \frac{\partial}{\partial y} q(y, t) = 0,$$

$$2 \left(\frac{\partial}{\partial y} v(y, t) \right) y + \left(\frac{\partial^2}{\partial y^2} v(y, t) \right) y^2 + a^2 \frac{\partial^2}{\partial t^2} v(y, t) - \mu^2 v(y, t) a^2 = 0,$$

$$2 \left(\frac{\partial}{\partial y} q(y, t) \right) y - 2 \left(\frac{\partial}{\partial t} v(y, t) \right) y + \left(\frac{\partial^2}{\partial y^2} q(y, t) \right) y^2 + \left(\frac{\partial^2}{\partial t^2} q(y, t) \right) a^2 - \mu^2 q(y, t) a^2 = 0.$$

It is equivalent to the following non homogeneous equation

$$- \left(\frac{\partial^2}{\partial y^2} \Phi(y, t) \right) y^2 - \left(\frac{\partial^2}{\partial t^2} \Phi(y, t) \right) a^2 + \mu^2 \Phi(y, t) a^2 + \epsilon = 0, \quad (41)$$

where

$$q(y, t) = \frac{\partial \Phi(y, t)}{\partial t}, v(y, t) = \frac{\partial \Phi(y, t)}{\partial y}$$

and ϵ is parameter.

The simplest solution of homogeneous equation can be presented in form

$$\Phi(y, t) = F_1(y) F_2(t), \quad (42)$$

where

$$\begin{aligned} \frac{d^2}{dt^2} F_2(t) &= -\frac{c_1 F_2(t)}{a^2} + \mu^2 F_2(t), \\ \frac{d^2}{dy^2} F_1(y) &= \frac{c_1 F_1(y)}{y^2}, \end{aligned} \quad (43)$$

and c_1 is the parameter.

The second equation from (43) has the form

$$\frac{d^2}{dy^2} F_1(y) - \frac{c_1 F_1(y)}{y^2} = 0$$

and its solutions are defined by the relations

$$\frac{F_1}{\sqrt{y}} = C_1 \cos(b \ln(y)) + C_2 \sin(b \ln(y)), \quad b^2 = -c_1 - \frac{1}{4} > 0,$$

$$\frac{F_1}{\sqrt{y}} = C_1 x^b + C_2 x^{-b}, \quad b^2 = \frac{1}{4} + c_1 > 0,$$

$$\frac{F_1}{\sqrt{y}} = C_1 + C_2 \ln(y), \quad c_1 = -\frac{1}{4},$$

which from the parameter c_1 are dependent.

The solutions of the first equation of the system (43) are

$$F_2(t) = C_3 \sin(1/4 \frac{\sqrt{2} \sqrt{c_1 - 8\mu^2 a^2} t}{a}) + C_4 \cos(1/4 \frac{\sqrt{2} \sqrt{c_1 - 8\mu^2 a^2} t}{a}).$$

In result the general solution of the equation (41) can be constructed with the help of solutions $F_1(y)$ and $F_2(t)$.

So depend upon the choice of c_1 the spectrum of manifold and the solutions of the equation (41) will be various.

The problem of solutions of the system (40) in more general case of the 1-form $\omega = A_i(x, y, z) dx^i$ requires more detail consideration.

6 Spatial metric of the four-dimensional Gödel space-time

The spatial metric of any four-dimensional metric

$$^4ds^2 = g_{\alpha\beta}dx^\alpha dx^\beta + 2g_{0\alpha}dx^0 dx^\alpha + g_{00}dx^0 dx^0$$

has the form

$$^3dl^2 = \gamma_{\alpha\beta}dx^\alpha dx^\beta,$$

where

$$\gamma_{\alpha\beta} = -g_{\alpha\beta} + \frac{g_{0\alpha}g_{0\beta}}{g_{00}}.$$

a three-dimensional tensor determining the properties of the space.

In the case of the Gödel space-time the spatial three-dimensional metric has the form

$$-^3dl^2 = \frac{a^2}{y^2}(dx^2 + dy^2) + dz^2. \quad (44)$$

Three-dimensional space with the metric (44) belongs to the one of the eight types of the W.Thurston geometries and has a diverse global properties.

In particular it admits the surfaces bundle.

As example we consider the translation surfaces of the space (44).

They defined by the system of equations for coordinates $x(u, v)$, $y(u, v)$ and $z(u, v)$

$$\begin{aligned} \frac{\left(\frac{\partial^2}{\partial u \partial v}x(u, v)\right) y(u, v) - \left(\frac{\partial}{\partial u}x(u, v)\right) \frac{\partial}{\partial v}y(u, v) - \left(\frac{\partial}{\partial u}y(u, v)\right) \frac{\partial}{\partial v}x(u, v)}{y(u, v)} &= 0, \\ \frac{\left(\frac{\partial^2}{\partial u \partial v}y(u, v)\right) y(u, v) + \left(\frac{\partial}{\partial u}x(u, v)\right) \frac{\partial}{\partial v}x(u, v) - \left(\frac{\partial}{\partial u}y(u, v)\right) \frac{\partial}{\partial v}y(u, v)}{y(u, v)} &= 0, \\ \frac{\partial^2}{\partial u \partial v}z(u, v) &= 0. \end{aligned} \quad (45)$$

The simplest solutions of these equations are in form

$$y(u, v) = 1/2 \left(1 + \left(\left(\frac{u}{v} \right)^{C_1} \right)^2 (e^{C_2 C_1})^{-2} \right) e^{C_2 C_1} \left(\left(\frac{u}{v} \right)^{C_1} \right)^{-1} C_1^{-1},$$

$$x(u, v) = \ln(u) + \ln(v),$$

and

$$z(u, v) = A(u) + B(v),$$

where $A(v)$ and $B(v)$ are arbitrary functions and C_1, C_2 are the parameters.

In particular case $C_1 = 1$, $C_2 = 0$ one get

$$y(u, v) = 1/2 \frac{u^2 + v^2}{uv},$$

$$x(u, v) = \ln(uv).$$

From here we find

$$\begin{aligned} v &= \sqrt{ye^x + e^x \sqrt{y^2 - 1}}, \\ u &= \frac{e^x}{\sqrt{ye^x + e^x \sqrt{y^2 - 1}}} \end{aligned}$$

and

$$z(x, y) = A(\sqrt{ye^x + e^x\sqrt{y^2 - 1}}) + B\left(\frac{e^x}{\sqrt{ye^x + e^x\sqrt{y^2 - 1}}}\right)$$

with arbitrary functions $A(u), B(v)$.

The properties of surfaces are dependent from the choice of the functions A and B .

Remark 3 From the system (45) we find the relations

$$\begin{aligned} \left(\frac{\partial}{\partial v}x(u, v)\right)^2 - \frac{e^{2z(u, v)}}{v^2} + \left(\frac{\partial}{\partial v}z(u, v)\right)^2 e^{2z(u, v)} &= 0, \\ \left(\frac{\partial}{\partial u}x(u, v)\right)^2 - \frac{e^{2z(u, v)}}{u^2} + \left(\frac{\partial}{\partial u}z(u, v)\right)^2 e^{2z(u, v)} &= 0, \end{aligned}$$

where $z(u, v) = \ln(y(u, v))$.

This fact allow us to get one equation on variable $y(u, v)$ only.

$$\begin{aligned} &\left(-\sqrt{(y)^2 - \left(\frac{\partial}{\partial v}y\right)^2} v^2 v \left(\frac{\partial}{\partial u}y\right) u^2 + \sqrt{(y)^2 - \left(\frac{\partial}{\partial u}y\right)^2} u^2 u \left(\frac{\partial}{\partial v}y\right) v^2\right) \frac{\partial^2}{\partial u \partial v} y + \\ &+ \sqrt{(y(u, v))^2 - \left(\frac{\partial}{\partial v}y\right)^2} v^2 v \left(\frac{\partial}{\partial v}y\right) y - \sqrt{(y)^2 - \left(\frac{\partial}{\partial u}y\right)^2} u^2 u \left(\frac{\partial}{\partial u}y\right) y = 0. \end{aligned}$$

References

- [1] Paterson E.M. and Walker A.G., Riemann extensions, *Quart.J.Math.Oxford*, 1952, v.3,19–28.
- [2] Dryuma V., Applications of Riemannian and Einstein-Weyl Geometry in the theory of second order ordinary differential equations, *Theoretical and Mathematical Physics*, 2001, v.128, N 1, 845–855.
- [3] Dryuma V., The Riemann Extension in theory of differential equations and their applications, *Matematicheskaya fizika, analiz, geometriya*, 2003, v.10, No.3,1–19.
- [4] Dryuma V., The Invariants, the Riemann and Einstein-Weyl geometry in theory of ODE's and all that, *A.B.Shabat et al. (eds.) New Trends in Integrability and Partial Solvability* 115–156, 2004, Kluwer Academic Publishers. ArXiv: nlin: SI/0303023, 11 March , 2003, 1–37.
- [5] Dryuma V., The Riemann Extension of the Schwarzschild space-time, *Buletinul A.S. M. Matematica*, Nr.3(43), 2003, p.92–103.
- [6] Dryuma V. On geodesical extension of the Schwarzschild space-time, *Proceedings of Institute of Mathematics of NAS of Ukraine*, Kiev, v.50, 2004, p.1294–1300.
- [7] Dryuma V. On the Riemann extension of the space-time with vanishing curvature invariants, <http://www.math.kth.se/4ecm/abstract/14.27.pdf>, Sweden, Stockholm, June 27–July 2, ECM'2004.
- [8] Dryuma V., The Riemann Extension of the Minkowsky space-time in rotating coordinate system, *ArXiv: gr-qc/0505122*, v1, 24 May 2005, p.1–10.
- [9] Dryuma V., On theory of the spaces of constant curvature, *ArXiv: math.DG/0505376*, 18 May 2005, p.1–12.
- [10] Dryuma V., Riemann extensions in theory of the first order systems of differential equations, *ArXiv:math.DG/0510526*, 25 October 2005, p.1–21.

- [11] Dryuma., From the Number theory to the Riemann geometry, *Scientific Papers International Conference "Differential Equations and Computer Algebra Systems (DE and CAS'2005)"*, Minsk, 2005, BGPU, Part 1, 14-15.
- [12] Hawking S.W., Ellis G., The Large Scale Structure of Space-Time, Moskow, 1977, (in russian).
- [13] Dautcourt G., Abdel-Megied M., Revisiting the Light Cone of the Gödel Universe, *ArXiv:gr-qc/0511015*, 3 Nov 2005, p.1-23.
- [14] Jackiw R., A Pure Cotton Kink in a Funny Place, *ArXiv: math-ph/0403044*, v.2, 21 July, 2004.
- [15] Eisenhart L.P., Riemannian geometry, Moskow, 1948, (in russian).
- [16] Elianu I.P., Cercetari asupra sistemelor de ecuatii lineare cu derivate partiale de tip Laplace, *Studii si cercetari matematice*, v.IV, Nr. 1-2, 1953, p.155-196.